

## Bax Recombinant antibody

**Cat:** B02004R

**Company:** HaoKebio

**Uniprot ID:** Q07812

**Applications:** IHC:1:500-1:2000

**Organism:** Rabbit

IHC-Polymer:1:2000-1:8000

**Species reactivity:** Human Mouse Rat

IHC-TSA:1:2500-1:10000

IF:1:50-1:100

WB:1:500-1:1000

### Background:

BAX (also known as BCL2 Associated X, Bcl-2-Like Protein 4, Bcl2-L-4, BCL2L4) is a member of the BCL2 family of proteins that play a key role in the regulation of apoptosis in higher eukaryotes. BAX comprises 4 Bcl-2 homology domains (BH1-BH4) and a C-terminal transmembrane domain. In healthy mammalian cells, BAX is localized to the cytoplasm through its interaction with the anti-apoptotic Bcl-2 family members BCL2L1/Bcl-xL. In response to apoptotic stimuli, however, BAX undergoes a conformational change that causes it to translocate to the outer mitochondrial membrane where it initiates the mitochondrial pathway of apoptosis via two potential mechanisms. Firstly, upon translocation to the outer mitochondrial membrane, BAX interacts with the mitochondrial voltage-dependent anion channel (VDAC) leading to the opening of the channel, loss of membrane potential, and the release of cytochrome c from the mitochondrion. The release of cytochrome C into the cytoplasm leads to the activation of Caspase3, initiating apoptosis. Secondly, activated BAX forms homodimers, which then assemble into oligomers on the mitochondrial outer membrane to create pores that permeabilize the mitochondrion leading to the release of cytochrome C.

### Protein full name:

BCL2-associated X protein

### Synonyms:

Apoptosis regulator BAX, Bcl2-L-4, Bcl-2-like protein 4

### Immunogen:

Recombinant protein

### Isotype:

IgG

### Subcellular location:

Cytoplasm

### Purity:

Affinity purification

### Form:

Liquid

### Storage Buffer:

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.


### Storage:

Store at -20 °C for one year.


### Experimental procedure:

Antigen retrieval: Citrate buffer (pH 6.0), Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

### Images:



Sample: Mouse tumor, 4% PFA 12-24h



Sample: Mouse tumor, 4% PFA 12-24h

**Source of Reagents:**

发表[中文论文]请标注:Bax(B02004R)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:Bax(B02004R) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.