

Species reactivity: Mouse

CD8 Recombinant antibody

Cat: B03042R Company: HaoKebio

Uniprot ID:P01731 Applications: IHC:1:200-1:2000

Organism: Rabbit IHC-Polymer: 1:800-1:8000

IHC-TSA:1:1000-1:10000

IF:1:50

WB:1:1000

Background:

CD8 is a transmembrane glycoprotein that is pre dominantly expressed on the surface of cytotoxic T cells, and can also be found on natural killer c ells, cortical thymocytes, and dendritic cells. CD 8 is composed of two disulfide-linked chains and can be present as a homodimer of CD8 a or as a heterodimer of CD8 α and CD8 β . The majority of class I-restricted T cells express mostly the C D8 α β heterodimer while CD8 α α homodimer s alone have been found on some gut intraepithel ial T cells, on some T cell receptor (TCR) γ δ T cells and on NK cells. CD8 acts as a co-recept or that binds to MHC class-I and participates in c ytotoxic T cell activation (PMID: 8499079). Duri ng T cell development, CD8 is required for positi ve selection of CD4-/CD8+ T cells.

Protein full name:

CD8 antigen, alpha chain

Synonyms:

CD8, CD8 antigen, alpha chain, CD8 α , Ly 2, L y 35

Immunogen:

Recombinant protein

Isotype:

IgG

Subcellular location:

Membrane

Purity:

Affinity purification

Form:

Liquid

Storage Buffer:

PBS with 0.02%sodium azide,100 μ g/ml BSA and 50% glyce rol.


Storage:

Store at -20 °C for one year.


Experimental procedure:

Antigen retrieval: Citrate buffer (pH 6.0), Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: P oly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

Images:




Sample: Mouse tumor, 4% PFA 12-24h.



Sample: Mouse tumor, 4% PFA 12-24h.

Source of Reagents:





发表[中文论文]请标注:CD8(B03042R)由杭州 浩克生物技术有限公司提供;

发表[英文论文]请标注:CD8(B03042R) were kin dly provided by Hangzhou Haoke Biotechnolog y Co., Ltd.