

NF-κB p65 Recombinant antibody

Cat:B14003R

Company: HaoKebio

Uniprot ID:Q04206

Applications: IHC:1:1000-1:5000

Organism:Rabbit

IHC-Polymer:1:4000-1:20000

Species reactivity:Human Mouse

IHC-TSA:1:5000-1:25000

IF:1:50-1:250

WB:1:1000-1:10000

Background:

Nuclear factor κB (NF-κB) is a sequence-specific DNA-binding protein complex which regulates the expression of viral genomes, including the human immunodeficiency virus, and a variety of cellular genes, particularly those involved in immune and inflammatory responses. The members of the NF-κB family in mammalian cells include the proto-oncogene c-Rel, p50/p105 (NFκB1), p65 (RelA), p52/p100 (NFκB2), and RelB. All of these proteins share a conserved 300-amino acid region known as the Rel homology domain which is responsible for DNA binding, dimerization, and nuclear translocation of NF-κB. The p65 subunit is a major component of NF-κB complexes and is responsible for trans-activation. NF-κB heterodimeric p65-p50 and p65-c-Rel complexes are transcriptional activators. The NF-κB p65-p65 complex appears to be involved in invasion-mediated activation of IL-8 expression. The inhibitory effect of IκB upon NF-κB in the cytoplasm is exerted primarily through the interaction with p65. p65 shows a weak DNA-binding site which could contribute directly to DNA binding in the NF-κB complex. It associates with chromatin at the NF-κB promoter region via association with DDX1. This antibody is a rabbit polyclonal antibody raised against residues near the N terminus of human RELA.

Protein full name:

v-rel reticuloendotheliosis viral oncogene homolog A (avian)

Synonyms:

NF κB, p65, p65; RELA, RELA, 4C7

Immunogen:

Recombinant protein

Isotype:

IgG

Subcellular location:

Cytoplasm, Nucleus

Purity:

Affinity purification

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 100 μg/ml BSA and 50% glycerol.


Storage:

Store at -20 °C for one year.

Experimental procedure:

Antigen retrieval: Citrate buffer (pH 6.0), Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

Images:



Sample: Mouse testicle, 4% PFA 12-24h

Source of Reagents:

发表[中文论文]请标注:NF- κ B p65(B14003R)由
杭州浩克生物技术有限公司提供;
发表[英文论文]请标注:NF- κ B p65(B14003R) w
ere kindly provided by Hangzhou Haoke Biotech
nology Co., Ltd.